

NEET: 2022

Candidate's Signature

PCB Test: 1

Time: 03 Hours

Invigilator's Signature

Ques	stion Booklet Version]	Roll N	umbe	r			9	Question Booklet Sr.	No.
11	(Write this number on your Answer Sheet)	0									
This is t	to certify that, the entries of R	CC-2022	2 Roll N	o. and A	Answer	Sheet 1	No. hav	e b	eer	n correctly written and ve	rified
			S	olut	tion	S					

NTA UPDATED QUESTION PAPER PATTERN

Sr. No.	Subject(s)	Section(s)	No. Of Question(s)	Mark(s)* *(Each Question Carries 04 (Four Marks))	Type Of Question(s)
	PHYSICS	SECTION A	35	140	
1.	11110100	SECTION B	15	40	
	CHEMISTRY	SECTION A	35	140	MCQ
2.	OTILIMIOTICI	SECTION B	15	40	(Multiple
	BOTONY	SECTION A	35	140	Choice Questions)
3.	20.0	SECTION B	15	40	Questions
	ZOOLOGY	SECTION A	35	140	
4.		SECTION B	15	40	
		TOTAL MARKS		720	

Note: ■ Correct option marked will be given (4) Marks and incorrect option marked will be minus one (-1) mark. Unattempted/Unonswered Questions will be given no marks.

■ Section B will have 15 questions, out of these 15 Questions, candidates can choose to attempt any10 Questions.

• Test Syllabus •

Physics: (11th + 12th) Complete Syllabus

Chemistry: (11th + 12th) Complete Syllabus

Biology: (11th + 12th) Complete Syllabus

Section 'A': Physics

RCC** RCC** RCC** RCC**

RCC** RCC** RCC** RCC** RCC** RCC** RCC** RCC**

RCC**

RCC** RCC**

RCC**

Section 'A'

1. A car is moving on a curved path with banking angle θ , the coefficient of friction between tyres of car and road is μ_c . The maximum safe velocity of car is


1)
$$\sqrt{gR^2 \left(\frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}\right)}$$
 2) $\sqrt{gR \left(\frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}\right)}$

$$2) \sqrt{gR\left(\frac{\mu_s + \tan\theta}{1 - \mu_s \tan\theta}\right)}$$

3)
$$\sqrt{\frac{g}{R} \left(\frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta} \right)}$$

3)
$$\sqrt{\frac{g}{R} \left(\frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta} \right)}$$
 4) $\sqrt{\frac{g}{R^2} \left(\frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta} \right)}$

Sol. (2):





$$F_{cp} = N \sin\theta + f \cos\theta$$
 (1)

$$Mg + f \sin\theta = N \cos\theta$$
(2)

The electric potential at a distance $\frac{R}{2}$ from the centre of a conducting sphere of charge Q & radius R will be




$$2) \ \frac{Q}{8\pi \in_0 R}$$

$$3) \ \frac{Q}{4\pi \in R}$$

$$4) \ \frac{Q}{2\pi \in_0 R}$$


Sol. (3):
$$V_{\text{(inside spher)}} = V_{\text{surface}}$$

Transfer characteristics [(output voltage (V_a)) vs input voltage (V)] for a base biased transistor in CE configuration is as shown in the figure. For using transistor as a switch, it is used.




- 1) in region II
- 2) in region I
- 3) in region III
- 4) both in region (I) & (III)

Sol. (4): As a switch In saturation (on switch)→ In cut off (off switch)



Three voltmeter A, B and C having resistances R and 3R



respectively are connected as shown in the figure. When some potential difference is applied between X and Y, the respective voltmeter readings are $V_{\scriptscriptstyle A}$, $V_{\scriptscriptstyle B}$ and $V_{\scriptscriptstyle C}$. Then

1)
$$V_A \neq V_B = V_C$$
 2) $V_B \neq V_A = V_C$
3) $V_A = V_B \neq V_C$ 4) $V_A = V_B = V_C$

2)
$$V_{R} \neq V_{A} = V_{A}$$

$$3) V_A = V_B \neq V_C$$

$$4) V_A = V_B = V_C$$

Sol. (4): B & C are in
$$\|^{\ell}$$
 & $R_A = R_B \| R_C$

- A screen is placed 50 cm from a single slit, which is illuminated with 6000 Å light. If distance between the first and third minima in the diffraction pattern is 3.00 mm, what is the width of the slit?
 - 1) 0.1 mm
- 2) 0.2 mm
- 3) 0.3 mm
- 4) 0.4 mm
- Sol. (2): In case of diffraction at single slit, the position of minima is given by

$$d \sin \theta = n\lambda$$
 ...(1)

and for small
$$\theta$$
: $\sin \theta = \theta = (y/D)$...(2)

So, from Eqs. (1) and (2), we have:

$$d\left(\frac{y}{D}\right) = n\lambda, \quad i. e. \quad y = \frac{D}{d}(n\lambda)$$

So that,
$$y_3 - y_1 = \frac{D}{d}(3\lambda - \lambda) = \frac{D}{d}(2\lambda)$$

and hence, d =
$$\frac{0.50 \times (2 \times 6 \times 10^{-7})}{3 \times 10^{-3}}$$

$$= 2 \times 10^{-4} \text{m} = 0.2 \text{ mm}$$




- 6. Power delivered by the ac source to the circuit becomes maximum, when
 - 1) $\omega L = \omega C$
- 2) $\omega L = \frac{1}{\omega C}$
- 3) $\omega L = \left(\frac{1}{\omega C}\right)^2$
- 4) $\omega L = \sqrt{\omega C}$
- **Sol. (2):** at resonance power is max so $X_L = X_C$
- 7. Focal length of a magnifying glass is 12.5 cm. Ratio of maximum and minimum magnifying power is
 - 1) 2:3
- 2) 1:3
- 3) 3:1
- 4) 3:2

Sol. (4):
$$MP_{min} = \frac{D}{f} = \frac{25}{12.5} = 2$$

 $MP_{max} = \frac{D}{f} + 1 = \frac{25}{12.5} + 1 = 3$

8. Same current i = 2A is flowing in a wire frame as shown in figure. The frame is a combination of two equilateral triangles ACD and CDE of side 1m. It is placed in uniform magnetic field B = 4T acting perpendicular to the plane of frame. The magnitude of magnetic force acting on the frame is




- 1) 24 N
- 2) Zero
- 3) 16 N
- 4) 8 N

Sol. (1): F = IBL

- 9. Avalanche breakdown in a PN junction diode is due to
 - 1) Sudden shift to Fermi level
 - 2) Increase in the width of forbidden gap
 - 3) Sudden increase of impurity concentration
 - 4) Cumulative effect of increased electron collision and creation of added electron hole pairs
- Sol. (4): Conceptual

10. In the following circuit, the reading of the voltmeter will be (in volts)



- 1) 7.2
- 2) 4.8

3) 6

RCC** RCC**

RCC**

RCC** RCC**

RCC**

RCC** RCC** RCC**

RCC** RCC**

RCC**

RCC** RCC**

- 4) 4
- **Sol. (1):** voltage on 40 cm = 4.8 V

Potential gradiant =
$$\frac{4.8}{40 \text{cm}}$$

voltage on 60 cm so =
$$\frac{4.8}{40} \times 60 = 7.2 \text{ volt.}$$

$$\Rightarrow \frac{5 \times 6 \times 10^{-7} \times 150 \times 10^{-2}}{0.5 \times 10^{-3}}$$
= 9 mm

- 11. Two identical metal plates show photoelectric effect by a light of wavelength λ_A falling on plate A and λ_B on plate B ($\lambda_A = 2\lambda_B$). The maximum kinetic energy is
 - $1) \ 2K_A = K_B$
- 2) $K_A < K_B / 2$
- 3) $K_A = 2K_B$
- 4) $K_A > K_B / 2$

Sol. (2):
$$\frac{hc}{\lambda_A} = K_A + W \Rightarrow \frac{hc}{2\lambda_B} = K_A + W$$

$$\Rightarrow \frac{hc}{\lambda_B} = 2K_A + 2W \dots (1)$$
and $\frac{hc}{\lambda_B} = K_B + W \dots (2)$

by (1) and (2)


$$2K_A + 2W = K_B + W \Rightarrow \frac{K_B}{2} - K_A = \frac{W}{2}$$

- 12. The half-life of a sample of a radioactive substance is 1 hour. If 8×10^{10} atoms are present at t = 0, then the number of atoms decayed in the duration t = 2 hour to t = 4 hour will be
 - 1) 2×10^{10}
- 2) 1.5×10^{10}
- 3) zero
- 4) Infinity

Sol. (2):
$$N = \frac{N_0}{2^2} - \frac{N_0}{2^4}$$




13. For what value of R in the circuit as shown current passing through 4 Ω resistance will be zero



- 1) 1 Ω
- $2) 2 \Omega$
- $3) 3 \Omega$
- 4) 4 Ω

Sol. (1):




 $6 = 10 - 2I \Rightarrow I = 2$ Amp.


Also 6 = 4 + IR

 $6 = 4 + 2(R) \implies R = 1\Omega$

14. A 14 kg block is hanged using a system of pulleys as shown in figure. Tension in string connecting celling and topmost pulley is



- 1) 17.5 N
- 2) 70 N
- 3) 140 N
- 4) 280 N
- Sol. (3): /////////




Therefore, 4T + 2T + T + T = 14g

$$8T = 140$$

Tension in string connecting ceiling and topmost pulley

15. In the circuit shown, when the switch is closed, the capacitor is charged with time constant τ_1 and when switch is open, then capacitor discharge with time constant τ, then $\tau_1 / \tau_2 is$



1) 1

RCC** RCC**

RCC** RCC** RCC** RCC** RCC**

RCC** RCC** RCC** RCC**

2) 1/2

3) 2

4) 1/4

Sol. (2):
$$\tau_1 = RC$$
 and $\tau = 2RC$ $\Rightarrow \frac{\tau_1}{\tau_2} = \frac{1}{2}$

- 16. A solid cylinder of mass M and radius R rolls down an inclined plane of height h. The angular velocity of the cylinder when it reaches the bottom of the plane is

 - 1) $\frac{1}{R}\sqrt{\frac{gh}{2}}$ 2) $\frac{2}{R}\sqrt{\frac{gh}{3}}$
 - 3) $\frac{1}{R}\sqrt{\frac{2gh}{3}}$ 4) $\frac{3}{R}\sqrt{\frac{2gh}{2}}$

Sol. (2):
$$I = \frac{1}{2}MR^2$$

When the cylinder rails down the inclined plane from a vertical height h its potential energy Mgh partly convened into translations kinetic energy $\frac{1}{2}Mv^2$, and partly into rotational

kinetic energy $\frac{1}{2}I\omega^2$. Therefore,

$$Mgh = \frac{1}{2}Mv^2 + \frac{1}{2}I\omega^2$$

$$= \frac{1}{2}Mv^2 + \frac{1}{2}\left(\frac{1}{2}MR^2\right)\omega^2$$
$$= \frac{1}{2}M(R\omega)^2 + \frac{1}{2}MR^2\omega^2 \quad (\because R\omega)$$


$$Mgh = \frac{3}{4}MR^2\omega^2$$

$$\omega = \sqrt{\frac{4gh}{3R^2}} = \frac{2}{R}\sqrt{\frac{gh}{3}}$$

ACC** RCC** RCC** RCC** RCC** RCC**




17. A man is crossing a river flowing with velocity of 5 m/s. He reaches at points B



directly across at a distance of 60m in 5 sec. His velocity in still water should be

- 1) 12 m/s
- 2) 13 m/s
- 3) 5 m/s
- 4) 10 m/s

Sol. (2):



$$u = \frac{60}{5} = 12 \text{ m/s}, v = 5 \text{ m/s}$$

 $\sqrt{v^2 + u^2} = \sqrt{5^2 + 12^2} = 13 \text{ m/s}$


- 18. If the density of the earth is doubled keeping its radius constant, then acceleration due to gravity will be $(g = 9.8 \text{ m/sec}^2)$
 - 1) 19.6 m/s²
- 2) 9.8 m/s²
- 3) 4.9 m/s^2
- 4) 2.45 m/s^2

Sol. (1):
$$g = \frac{4}{3}\pi R \rho g$$
 $g \propto \rho$

- 19. Mars has a diameter of approximately 0.5 of that of earth and mass of 0.1 of that of earth. The surface gravitational field strength on mars as compared to that on earth is greater by a factor of
 - 1) 0.1
- 2) 0.2
- 3) 2.0
- 4) 0.4

Sol. (4):
$$g_E = \frac{GM_E}{R_E^2}$$
, $g_M = \frac{GM_M}{R_M^2}$
 $g_m = \frac{G(0.1M_E)}{(0.5R_E)^2} = 0.4g_E$

20. A block of mass 1 kg is pushed up surface inclined to horizontal at angle of 30° by a force of 10 N parallel to the



inclined surface as shown in the figure. The coefficient of friction between block and the incline is 0.1. If the block is pushed up by 10 m along the inclined, the work done against force of friction is

- 1) 8.7 J
- 2) 10.7 J
- 3) 7.8 J
- 4) 12.7 J

Sol. (1): Work against friction is

$$\begin{split} W_f &= fd = \mu Nd = \mu mgd \, cos\theta \, (:: N = mgcos\theta) \\ &= 0.1 \times 1 \, kg \times 10 \, ms^{-2} \times cos \, 30^\circ \times 10m \\ &= 8.7 \, J \end{split}$$

- 21. A particle is moving with the velocity $v = (4t^3 + 3t^2 - 1)$ m/s. The displacement of particle in time t = 1 sec to t = 2sec will be

RCC**]

RCC** RCC**

RCC**

RCC** RCC**

RCC** RCC**

RCC** RCC**

RCC** RCC**


RCC**

RCC** RCC**

- 2) 17 m
- 3) 13 m
- 4) 9 m

Sol. (1):
$$\int ds = \int_{1}^{2} v dt = \int_{1}^{2} (4t^{3} + 3t^{2} - 1) dt$$
$$s = (t^{4} + t^{3} - t)_{1}^{2} = 21 \text{ m}$$

22. In the adjoining figure, the tension in the string connecting A and B is



- 1) g 2) $\frac{g}{9}$ 3) $\frac{8g}{9}$ 4) $\frac{10g}{9}$

Sol. (4):
$$a = \frac{5g - 4g}{9} = \frac{g}{9}$$

$$T_{AB} - 1g = 1 \times \frac{g}{9} \implies T_{AB} = \frac{10 g}{9}$$

- 23. If a body is executing simple harmonic motion, then
 - 1) at extreme position, the total energy must be
 - 2) at equilibrium position, the total energy is in the form of only potential energy
 - 3) at equilibrium position, the total energy is in the form of only kinetic energy
 - 4) at extreme position, the total energy is only potential energy
- **Sol. (4):** At extreme position kinetic energy is zero, total energy in the form of potential energy. At equilibrium kinetic energy is maximum but potential energy may or may not be zero. (Example vertical oscillations of spring)



- 24. The angle between two vectors given by $6\hat{i} + 6\hat{j} - 3\hat{k}$ and $7\hat{i} + 4\hat{j} + 4\hat{k}$ is

 - 1) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ 2) $\cos^{-1}\left(\frac{5}{\sqrt{3}}\right)$
 - 3) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$ 4) $\sin^{-1}\left(\frac{\sqrt{5}}{3}\right)$
- **Sol. (4)**: $\cos \theta = \frac{\vec{A} \cdot \vec{B}}{AB} = \frac{(6\hat{i} + 6\hat{j} 3\hat{k}) \cdot (7\hat{i} + 4\hat{j} + 4\hat{k})}{\sqrt{6^2 + 6^2 + 3^2} \sqrt{7^2 + 4^2 + 4^2}}$ $\cos \theta = \frac{42 + 24 - 12}{9 \times 9} = \frac{54}{9 \times 9} = \frac{2}{3}$ $\sin\theta = \sqrt{1 - \cos^2\theta} = \frac{\sqrt{5}}{2}$
- 25. We have two spheres one of which is hollow and the other solid. They have identical masses and moment of inertia about their respectively diameters. The ratio of their radius is given by
 - 1) 5:7
- 3) $\sqrt{3} \cdot \sqrt{5}$
- 4) $\sqrt{3}:\sqrt{7}$
- **Sol. (3):** $I_1 = I_2 \Rightarrow \frac{2}{3}MR_1^2 = \frac{2}{5}MR_2^2$ $\Rightarrow 5R_1^2 = 3R_2^2 \Rightarrow \frac{R_1}{R_2} = \sqrt{\frac{3}{5}}$
- 26. If $a = 8 \pm 0.08$ and $b = 6 \pm 0.06$, Let x = a + b, y = a - b, $z = a \times b$. The correct order of % error in x,y and z
 - 1) x = y < z
- 2) x = y > z
- 3) x < z < y
- 4) x > z < u
- **Sol. (3):** $a + b = 14 \pm 0.14 \Rightarrow \% \text{ error} = \frac{0.14}{12} \times 100 = 1\%$ $a - b = 2 \pm 0.14 \Rightarrow \% \text{ error} = \frac{0.14}{2} \times 100 = 7\%$ $a \times b = 48 \pm 0.96 \Rightarrow \% \text{ error} = \frac{20.96}{48} \times 100 = 2\%$ so, order of % error x < z < y
- 27. A mass of 100 gm is tied to one end of a string 2 m long. The body is revolving in a horizontal circle making a maximum of 200 revolutions per min. The other end of the string is fixed at the centre of the circle of revolution. The maximum tension that the string can bear is (approximately)
 - 1) 8.76 N
- 2) 8.94 N
- 3) 87.6 N
- 4) 896 N
- Sol. (3): $m\omega_{max}^2 r = T_{max}$

- 28. On heating one end of a rod, the temperature of whole rod will be uniform when
 - 1) K = 1
- 2) K = 0
- 3) K = 100
- 4) $K = \infty$
- **Sol.** (4): For uniform temperature on heating the rod, K should be infinite
- 29. Energy levels A, B and C of a certain atom correspond to increasing values of energy i.e. $E_A < E_B < E_C$. If λ_1 , λ_2 and λ_3 are wave lengths of radiations corresponding to transitions C to B, B to A and C to A respectively, which of the following relations is correct
 - 1) $\lambda_2 = \lambda_1 + \lambda_2$

RCC** RCC**


- 2) $\lambda_1 + \lambda_2 + \lambda_3 = 0$
- 3) $\lambda_3^2 = \lambda_1^2 + \lambda_2^2$ 4) $\lambda_3 = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$
- **Sol.** (4): $E_{C \rightarrow A} = E_{C \rightarrow B} + E_{B \rightarrow A}$

$$\frac{hc}{\lambda_3} = \frac{hc}{\lambda_1} + \frac{hc}{\lambda_2} = \lambda_3 = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$$

- 30. A uniform electric field of 20 N/C exists along the positive x-axis in space. The potential difference $(V_B - V_A)$ for the points A (4m, 2m) and B (6m, 5m) is
 - 1) $20\sqrt{13}$ volt
- 2) 40 volt
- 3) + 40 V
- 4) $-20\sqrt{13}$ volt
- Sol. (2): $\int_{\mathbf{E}} d\mathbf{v} = -\int_{\mathbf{E}} \vec{\mathbf{E}} \cdot d\vec{\mathbf{r}}$

$$V_B - V_A = -\int_{V_A}^{V_B} (20\hat{i}) \cdot (dxi) = -40 \text{ Volt}$$


31. Force at t = 3 sec is equal to



- 1) 1 N
- 2) -1 N
- 3) Zero
- 4) 3 N
- $|\mathbf{Sol.}|$ (2): F = slope of p t curve at t = 3, slope = -1.



32. Two identical particles having the same mass m and charges +q and -q separated by a distance d enter in a uniform magnetic field Bdirected perpendicular to paper inwards with speeds v_1 and v_2 as shown in figure. The particle will not collide if




- 1) $d > \frac{m}{Bq}(v_1 + v_2)$ 2) $d < \frac{m}{Bq}(v_1 + v_2)$
- 3) $d > \frac{2m}{Ba}(v_1 + v_2)$
- 4) $v_1 = v_2$
- Sol. (3):

for particle to not collide $(2r_1 + 2r_2) < d$

$$2\left(\frac{mv_1}{qB} + \frac{mv_2}{qB}\right) < d$$
$$d = \frac{2m}{qB}(v_1 + v_2)$$

33. The following configuration of gate is equivalent to



- 1) NAND
- 2) XOR
- 3) OR
- 4) None of these

- Sol. (2): $y = (A + B) \cdot (\overline{A \cdot B})$ $=(A+B) \cdot (\overline{A} + \overline{B})$ $= A\overline{A} + A\overline{B} + B\overline{A} + B \cdot \overline{B}$ RCC** 1 $= O + A\overline{B} + B\overline{A} + O = XOR$ Gate
 - 34. Two whistles A and B produce notes of frequencies 660 Hz and 596 Hz respectively. There is a listener at the mid-point of the line joining them. Both the whistle B and the listener start moving with speed 30 m/s away from whistle A. If speed of sound be 330 m/s, how many beats will be heard by the listener?
 - 1) 2

RCC** RCC**

RCC** RCC** RCC** RCC**

RCC** RCC** RCC** RCC**

RCC** RCC**

RCC** RCC**

RCC**

2) 4

3) 6

4) 8

Sol. (2):
$$fo_A=660 \text{ Hz}$$

$$A \qquad D \qquad Fo_B=596 \text{ Hz}$$

$$A \qquad D \qquad B \qquad 30 \text{ m/s}$$
(Listner)

Apparent frequency for whistle by 'A'

$$(f_{app})_A = fo_A \left(\frac{V - V_L}{V - V_S}\right)$$

= $660 \left(\frac{330 - 30}{330 - 0}\right) = 600 \text{ Hz}$

$$(f_{app})_{B} = f_{oB} \left(\frac{V - V_{L}}{V - V_{S}} \right)$$


= 596 $\left[\frac{330 - (-30)}{330 - (-30)} \right] = 596 \text{ Hz}$

Beat frequency $f_b = (f_{app})_{\Delta} - (f_{app})_{B}$ = 600 - 596 = 4 beats /sec

- 35. Which is the correct statement
 - 1) For an isothermal change PV = constant
 - 2) In an isothermal process the change in internal energy must be equal to the work done
 - 3) For an adiabatic change $\frac{P_2}{P_1} = \left(\frac{V_2}{V_1}\right)^{\prime}$, where λ is the ratio of specific heats
 - 4) In an adiabatic process work done must be equal to the heat entering the system
- **Sol.** (1): Since PV = RT and T =constant;
 - $\therefore PV = constant$

Section 'B'

36. Output Y of the given logic gate network is



- 1) $\overline{A} \cdot B = A \cdot \overline{B}$
- 2) $A.B = \overline{A}.\overline{B}$
- 3) $(\overline{A+B}) = \overline{A} \cdot \overline{B}$
- 4) None

Sol. (4):
$$Y = \frac{\overline{\overline{AB}} - \overline{\overline{AB}}}{\overline{AB}} = (\overline{AB}). (A\overline{B}) = 0$$

- 37. A 50-turn circular coil of radius 2.0 cm carrying a current of 5.0 A is rotated in a magnetic field of strength 0.20 T. In a particular position of the coil, the torque acting is half of the maximum torque. The angle between the magnetic field and the plane of the coil is
 - 1) 60°
- 2) 30°
- 3) 90°
- 4) 100°
- **Sol. (1) :** \angle between M & B is 30° so between B & plane of coil = 60°
- 38. In a Young's double slit experiment, slits are separated by 0.5 mm, and the screen is placed 150 cm away. A beam of light consisting of two wavelengths, 600 nm and 500 nm, is used to obtain interference fringes on the screen. The least distance from the common central maximum to the point where the bright fringes due to both the wavelengths coincide is
 - 1) 90 μm
- 2) 900 μm
- 3) 9 μm
- 4) 9 mm

Sol. (4):
$$\frac{n_1}{n_2} = \frac{\lambda_2}{\lambda_1} = \frac{500 \text{ nm}}{600 \text{ nm}}$$

$$n_1 = 5 \quad n_2 = 6$$

$$x = \frac{n_1 \lambda_1 D}{d}$$

$$\Rightarrow \frac{5 \times 6 \times 10^{-7} \times 150 \times 10^{-2}}{0.5 \times 10^{-3}}$$
= 9 mm

39. The electric field part of an electromagnetic wave in a medium is represented by E_z = 0;

$$E_{y} = 2.5 \frac{N}{C} \cos \left[\left(2\pi \times 10^{6} \frac{rad}{m} \right) t - \left(\pi \times 10^{-2} \frac{rad}{s} \right) x \right];$$

 $E_z = 0$. The wave is

RCC** RCC**

RCC** RCC**

RCC** RC

RCC**

RCC**

RCC** RCC**

RCC**

- 1) Moving along –*x* direction with frequency 10⁶ Hz and wave length 200 m
- 2) Moving along *y* direction with frequency $2\pi \times 10^6$ Hz and wave length 200 m
- 3) Moving along x direction with frequency 10^6 Hz and wave length 100 m
- 4) Moving along x direction with frequency 10^6 Hz and wave length 200 m


Sol. (4):
$$E_y = 2.5 \frac{N}{C} \cos[(2\pi \times 10^6 t) - \pi \times 10^{-2} x]$$

direction - X axis
wavelength $\lambda = \frac{2\pi}{k} = \frac{2\pi}{\pi \times 10^{-2}} = 200 \text{ m}$
 $f = \frac{\omega}{2\pi} = \frac{2\pi \times 10^6}{2\pi} = 10^6 \text{ Hz}$

- 40. The binding energy of deuteron ${}_{1}^{2}H$ is 1.112 MeV per nucleon and an α -particle ${}_{2}$ He 4 has a binding energy of 7.047 MeV per nucleon. Then in the fusion reaction , ${}_{1}^{2}H+{}_{1}^{2}H\rightarrow {}_{2}^{4}He+Q$, the energy Q released is
 - 1) 1 MeV
- 2) 11.9 MeV
- 3) 23.8 MeV
- 4) 931 MeV

Sol. (3):
$$Q = BE_{(prod)} - BE_{(React)}$$

41. A light beam is traveling from Region I to Region IV (Refer Figure). The refractive index in Regions I, II, III and IV are n_0 , $\frac{n_0}{2}$, $\frac{n_0}{6}$ and $\frac{n_0}{8}$, respectively. The angle of incidence θ for which the beam just misses entering Region IV in figure



- 1) $\sin^{-1}\left(\frac{3}{4}\right)$
- 2) $\sin^{-1}\left(\frac{1}{8}\right)$
- 3) $\sin^{-1}\left(\frac{1}{4}\right)$
- 4) $\sin^{-1}\left(\frac{1}{3}\right)$



Sol. (2):
$$n_0 \sin \theta = \frac{n_0}{8} \sin 90^\circ$$
 $\sin \theta = \frac{1}{8}$

42. A uniform rope of length L and mass m, hangs vertically from a rigid support. A block of mass m₂ is attached to the free end of the rope. A transverse pulse of wavelength λ_1 is produced at the lower end of the rope. The wavelength of the pulse when it reaches the top of the rope is λ_2 . The ratio λ_2 / λ_1 is RCC** RCC** RCC**

$$1) \sqrt{\frac{m_1}{m_2}}$$

2)
$$\sqrt{\frac{m_1 + m_2}{m_2}}$$

3)
$$\sqrt{\frac{m_2}{m_1}}$$

4)
$$\sqrt{\frac{m_1 + m_2}{m_1}}$$

Sol. (2):
$$V = \sqrt{\frac{T}{\mu}} \Rightarrow V_1 = \sqrt{\frac{m_2 g}{\mu}}$$

$$\Rightarrow V_2 = \sqrt{\frac{(m_1 + m_2)g}{\mu}}$$

$$V = f\lambda$$

$$\lambda \propto V$$

$$\frac{\lambda_2}{\lambda_1} = \sqrt{\frac{m_1 + m_2}{m_2}}$$

- 43. A body of uniform cross-sectional area floats in a liquid of density thrice its value. The portion of exposed height will be
 - 1) 2/3

2) 5/6

3) 1/6

- 4) 1/3
- **Sol. (1):** For floating body $W = Th = V_{in} \rho_L g$

$$\begin{split} &AH \; \rho_B g = (Ah) \times 3 \rho_B g \\ &\frac{h}{H} = \frac{1}{3} \\ &Exposed \; height \; ratio = \frac{H-h}{H} = \frac{3-1}{3} = \frac{2}{3} \end{split}$$


- 44. At a pressure of 24 × 10⁵ dyne cm⁻², the volume of O₂ is 10 litre and mass is 20 g. The rms velocity will be
 - 1) 800 ms⁻¹
- 2) 400 ms⁻¹
- 3) 600 ms⁻¹
- 4) Data is incomplete


Sol. (3):
$$C = \sqrt{\frac{3PV}{M}}$$

$$= \sqrt{\frac{3 \times 24 \times 10^5 \times 10 \times 100}{20}} \text{ cms}^{-1}$$

$$= 6 \times 10^4 \text{ cm s}^{-1} = 600 \text{ ms}^{-1}$$


45. A sound source, emitting sound of constant frequency, moves with a constant speed and crosses a stationary observer. The frequency (n) of sound heard by the observer is plotted against time (t). Which of the following graphs represents the correct variation





Source
When Approaching =
$$n \left[\frac{V}{V - V_s} \right]$$

When it's going away =
$$n \left[\frac{V}{V + V_s} \right]$$



- 46. A variable force, given by the 2-dimensional vector $\vec{F} = (3x^2\vec{i} + 4\vec{i})$, acts on a particle. The force is in newtons and x is in metres. What is the change in the kinetic energy of the particle as it moves from the point with coordinates (2, 3) to (3, 0)? (The coordinates are in metres)
- 2) Zero
- 4) +19 I


RCC** RCC**


RCC** RCC**


RCC** RCC** RCC**




- 47. Two light wave from coherent sources superimpose at point A with phase difference 0 & at point B with phase difference of π / 2. Calculate ratio of resultant intensities of point A and B
 - 1) 1:1
- 2) 2:1
- 3) 4:1
- 4) 1:4
- Sol. (2): $(I_{res})_A = I + I + 2\sqrt{I}\sqrt{I}\cos 0^{\circ}$ $(I_{res})_B = 2I$ $(I_{res})_A = 4I$ $(I_{res})_B = I + I + 2\sqrt{I}\sqrt{I}\cos 90^{\circ}$ $\frac{(I_{res})_B = 2I}{(I_{res})_A} = \frac{4I}{2I} = \frac{2}{1}$
- 48. In which of the following cases the contact force between A & B is maximum? ($m_A = m_B = 1 \text{ kg}$, g =10 ms⁻²)









Sol. (2): (1)
$$a = \frac{50}{2} = 25 \text{ m/sec}^2$$

B

 $\Rightarrow N = 1 \times a = 25 \text{ Newton}$

(2)

 $N = 40 \text{ Newton}$

 $N = 1 \times 35 = 35$ Newton

- 1) 12.42×10^{-21} J, 968 m/s
- 2) 8.78×10^{-21} J, 684 m/s
- 3) 6.21×10^{-21} J, 968 m/s
- 4) 12.42×10^{-21} J, 684 m/s

Sol. (4):
$$E = \frac{3NKT}{2}$$
, $V_{rms} = \sqrt{\frac{3KT}{m}}$
 $T_1 = 300 \text{ K}$, $T_2 = 600 \text{ K}$
 $\frac{E_2}{E_1} = \frac{T_2}{T_1} = 2 \Rightarrow E_2 = 2E_1$
 $E_2 = 2 \times 6.21 \times 10^{-21} = 12.42 \times 10^{-21} \text{ J}$
 $\frac{V_2}{V_1} = \sqrt{\frac{T_2}{T_1}} = \sqrt{2} \Rightarrow V_2 = 484 \times \sqrt{2}$

1) 1 gm

RCC** RCC**

RCC**

RCC** RCC**

RCC** RCC** RCC**

RCC** RCC**

RCC**

RCC** RCC**

RCC**

- 2) 10 kg
- 3) 100 kg
- 4) 100 gm

 $V_2 = 684 \text{ m/s}$

$$^{235}_{92}U \rightarrow 200 \text{ MeV}$$

in N_A Reaction $E = 200 \text{ MeV} \times N_A$

means energy emitted by 235 gram

$$_{92}$$
U $^{235} \rightarrow N_A \times 200 \text{ MeV}$

by m gram uranium
$$\rightarrow \frac{(N_A \times 200 MeV) \times m}{235}$$

If it is in one day then

$$\frac{\text{(N}_{A} \times 200 \text{MeV)m}}{235} = (0.95 \text{ Mw}) \times 1 \text{ day}$$

$$m = \frac{0.95 \text{MW} \times (24 \times 3600 \text{ sec}) \times 235}{\text{N}_{A} \times 200 \text{MeV}}$$

$$m = \frac{0.95 \times 10^6 \times 24 \times 3600 \times 235}{6 \times 10^{23} \times 200 \times 10^6 \times 1.6 \times 10^{-19}} \, gram$$
 on solving

$$m = 1 gram$$

Section 'B': Chemistry

RCC** RCC** RCC**

Section 'A'

- 51. Number of Oxygen atoms are maximum in
 - 1) 0.2 moles of BaCO₃
 - 2) 1 mole of H₃PO₄
 - 3) $0.5 \text{ moles of } C_6 H_{12} O_6$
 - 4) 0.75 mole of CO₂
- Sol. (2) 0.2 mole $BaCO_3 \equiv 0.6$ mole O atoms

 1 mole $H_3PO_4 \equiv 4$ mole O atoms

 0.5 mole $C_6H_1O_6 \equiv 3$ mole O atoms

 0.75 mole $CO_2 \equiv 1.5$ mole O
- 52. An element has 2 electrons in K shell, 8 electrons in L shell, 13 electrons in M shell and one electron in N shell. The element is
 - 1) Cr
- 2) Fe
- 3) V
- 4) Ti

Sol. (1)
$$= 2+8+13+1 = 24$$
 (C_{π})

- 53. The element with highest electron gain enthalpy will belong to
 - 1) Period 2, group 17
 - 2) Period 3, group 17
 - 3) Period 2, group 18
 - 4) Period 2, group 1

Sol. (2) It is Chlorine

- 54. During the change of O_2 to O_2^- , the incoming electron goes to the orbital
 - 1) σ*2pz
- 2) π2py
- 3) $\pi^{*}2px$
- 4) π2px
- Sol. (3) $O_2 = \sigma 1s^2 \sigma^* 1s^2 \sigma 2s^2 \sigma^* 2s^2$ $\sigma^2 P_z^2 \pi^2 P_z = \pi^2 P_y$ $\pi^* \frac{1}{2} P_z = \pi^* \frac{1}{2} P_y \sigma^* 2P_z$ So in the form of O^2 electron is added to
 either $\pi^* 2P_z = \sigma^* \pi^* 2P_y$,

- 55. The best method to separate the mixture of ortho and para nitrophenol (1:1) is
 - 1) Steam distillation 2) Crystallisation
 - 3) Vapourisation
- 4) Colour spectrum
- Sol. (1) Orthonitrophenol has Higher Vapour pressure due to intramolecular hydrogen bonding than paranitrophenol So can be separated by Steam distillation.
- 56. The correct order of increasing s-character (in percentage) in the hybrid orbitals of the following molecule/ion is
 - I. CO₃²⁻

II. XeF

III. I₃-

IV. NCl₃

V. BeCl,

- 1) II < III < IV < I < V
- 2) II < IV < III < V < I
- 3) III < II < I < V < IV
- 4) II < IV < III < I < V
- Sol. (1) $Becl_2 \Rightarrow SP \Rightarrow 50 \% S$ $CO_3^2 \Rightarrow SP^2 \Rightarrow 33.33\% S$ $NCl_3 \Rightarrow SP^3 \Rightarrow 25\% S$ $L_3^- \Rightarrow SP^3 d \Rightarrow 20\% S$ $XeF_4 \Rightarrow SP^3 d^2 \Rightarrow 16.3\% S$
- 57. At very high pressures, the compressibility factor of one mole of a gas is given by
 - 1) $1 + \frac{Pb}{RT}$
- 2) $\frac{Pb}{RT}$
- 3) $1-\frac{Pb}{RT}$
- 4) $1 \frac{b}{(VRT)}$
- Sol. (1) At very high pressure, the deviation so Z>1.



- 58. For a reaction, $\Delta H = -40 \text{ kJ}$ and $\Delta S = -50 \text{ J/K}$. At what temperature range will it change from spontaneous to non-spontaneous?
 - 1) 0.8 K to 1 K
- 2) 799 K to 800 K
- 3) 800 K to 801 K
- 4) 799 K to 801 K
- Sol. (4) Teq = $\frac{\Delta H}{\Delta S} = \frac{40 \times 10^3}{50}$ = 800 k

as both AH and As are -ve, rean is spontaneous at low T & nonspontaneous

at High T. Spontaneous at 799 K

of nonspontaneous at 801K.

59. For the reaction,

 $PCl_{5}(g) \rightleftharpoons PCl_{3}(g) + Cl_{5}(g)$

The forward reaction at constant temperature is favoured by :

- 1) Introducing an inert gas at constant volume
- 2) Introducing chlorine gas at constant volume
- 3) Introducing an inert gas at constant pressure
- 4) Introducing PCl₅ at constant pressure
- Sol. (3) When an inest gas is added at constant pressure, equilibrium Shifts in that direction where gaseous moles are greater.
- 60. A 50 ml solution of pH = 1 is mixed with a 50 ml solution of pH = 2 the pH of the mixture is
 - 1) 0.86
- 2) 1.26
- 3) 1.76
- 4) 2.26
- Sol. (2) Use $[H^{+}] = \frac{M_1V_1 + M_2V_2}{Vmix}$ = 55×10^{-3} PH = $-log(55 \times 10^{-3}) = 1.26$

- 61. Which of the following is not redox reaction:
 - 1) NaOH + HCl \rightarrow NaCl + H,O
 - 2) $2H_2 + O_2 \rightarrow H_2O_2$
 - 3) Na + $H_2O \rightarrow NaOH + 1/2H_2$
 - 4) $H_2 + Br_2 \rightarrow 1HBr$
- Sol. (1) Acid-Base neutralisation is nonredox reaction.
- 62. Select incorrect order

RCC** RCC** RCC**

- 1) Be(OH)₂ < Mg(OH)₂ < Ca(OH)₂ Solubility in H₂O
- 2) BeCO₃ < MgCO₃ < CaCO₃ Thermal stability
- 3) Cs < Rb < K < Na < Li Ease of formation of hydrides
- 4) CsH > RbH > KH > NaH Stability
- Sol. (4) Correct thermal stability
 Order of hydrides of
 aikali metals.
 LiH>NaH> KH> RbH> CsH.
- 63. Which comparison is not correct as indicated?
 - 1) OH>CH₃OH (acidic nature)
 - 2) \sim NH₂ > CH₃NH₂ (basic nature)

 - $4) \begin{picture}(200,0) \put(0,0){\line(0,0){0.5ex}} \put(0,0){\line(0,0){0$
- Sol. (2) Resonance $\propto \frac{1}{\text{Basic Nature}}$

(in

CH3-NH2

Resonance

No Resonance

The aniline, due to the presence of resonance, lone pain of electron, is not easily available boun protonation and so aniline is less basic than ey-Ny,



64. The most stable carbanion among the following is

$$\begin{array}{ccc} CH_2-CH_2^- & CH_2^- \\ \end{array}$$

$$\begin{array}{cccc} CH_2 - CH_2^- & CH_2^- \\ \end{array}$$

3)
$$\bigcirc$$
 CH₂ \bigcirc CH \bigcirc OCH₃ \bigcirc NO

65. The appropriate reagent for the following transformation is

- 1) Zn-Hg/HCl
- 2) H₂N-NH₂, KOH
- 3) Ni/H₂
- 4) NaBH₄

66. Incorrect match is

- 1) Permutit → Hydrated silicates of Na and Al
- 2) Calgon → Sodium Hexameta phosphate
- 3) BeH₂, MgH₂ \rightarrow Covalent polymeric hydride
- 4) Hydrolith → electron deficient hydride

Sol. (4) Hydrolith = CaH2 is ionic hydride.

- 67. A solid compound contains x, y, z atoms in a cubic lattice with x atoms occupying the corner, y atoms in the body centered positions and z atoms at the centres of faces of the unit cell. What is the empirical formula of the compound?
 - $1) XY_2Z_3$

RCC** RCC** RCC** RCC** RCC** RCC** RCC** RCC** RCC**

RCC** RCC** RCC** RCC** RCC**

- 2) XYZ₃
- 3) $X_{2}Y_{2}Z_{3}$
- 4) X_8YZ_6

Sol. (2)
$$\times = \frac{1}{8} \times 8 = 1$$

 $Y = 1$
 $Z = 6 \cdot \frac{1}{2} \times 6 = 3$
 $\times : Y : Z = 1 : 1 : 3 \Rightarrow \times YZ_3$

- 68. If sodium sulphate is considered to be completely dissociated into cations and anions in aqueous solution, the change in freezing point of water (ΔT_f), when 0.01 mol of sodium sulphate is dissolved in 1 kg of water is ($K_f = 1.86$ kg mol⁻¹)
 - 1) 0.0186 K
- 2) 0.0372 K
- 3) 0.0558 K
- 4) 0.0744 K

Sol. (3)
$$Na_2 so_4 \Rightarrow i=3$$

$$\Delta T_f = i K_f m$$

$$= 3 \times 1.86 \times 0.01$$

- 69. What would be the ratio of moles each of Ag⁺, Cu⁺², Fe⁺³ ions would be deposited by passage of same quantity of electricity through solutions of their salts
 - 1) 1:1:1
- 2) $1:\frac{1}{2}:\frac{1}{3}$
- 3) $\frac{1}{3}$: $\frac{1}{2}$: 1
- 4) 1:2:3



Sol. (2) Let charge passed = IF

$$Ag^{+} + e^{-} \xrightarrow{IF} Ag (Imole)$$

$$Cu^{2+} + 2e^{-} \xrightarrow{IF} Cu (\frac{1}{2} mole)$$

$$Fe^{3+} + 3e^{-} \xrightarrow{IF} Fe (\frac{1}{3} mole)$$

$$Ag^{+} : Cu^{2+} : Fe^{3+} = I : \frac{1}{2} : \frac{1}{3}$$

70. Given:

$$E^{\text{o}}_{\text{Ag}^{\text{+}}/\text{Ag}}$$
 = +0.80V , $E^{\text{o}}_{\text{Co}^{\text{2+}}/\text{Co}}$ = -0.28V

$$E^{\text{o}}_{\text{Cu}^{2^{\text{+}}}/\text{Cu}}$$
 = +0.34V , $E^{\text{o}}_{\text{Zn}^{2^{\text{+}}}/\text{Zn}}$ = -0.76V

Which metal will corrode fastest?

- 1) Ag
- 2) Cu
- 3) Co
- 4) Zn

Sol. (4) Reactivity of metal
$$\propto 1/E^{\circ}$$
As Zn is most reactive among given metal, it
Will Corrode fastest.

71. Which of the following does not affect the rate of reaction?

- 1) Amount of the reactants taken
- 2) Physical state of the reactants
- 3) ΔH of reaction
- 4) Size of the vessel

Sol. (3)
$$\Delta H$$
 do not affect the rate of reaction.

72. Electrolytic refining is used to purify which of the following metals?

- 1) Cu, Ag, Pb and Zn 2) Ge and Si
- 3) Zr and Ti
- 4) Zn and Hg


73. In the statements regarding P_4 molecule

- i) the oxidation state is zero
- ii) the covalency is 4
- iii) the P–P–P bond angle 60°

the correct combination is

- 1) only iii is correct 2) i and iii are correct
- 3) all are correct
- 4) i and ii are correct

Sol. (2) Covalency of P4 is 3



Each P form 3 covalent bonds. 50 covalency of each P=3.

74. Very pure N, can be obtained by

- 1) Thermal decomposition of ammonium dichromate
- 2) Treating aqueous solution of NH₄Cl and NaNO₂
- 3) Liquefaction and fractional distillation of liquid air
- 4) Thermal decomposition of sodium azide

75. Which of the following has peroxy linkage?

- 1) $H_2S_2O_3$
- 2) H₂SO₅
- 3) H₂S₂O₇
- 4) H₂S₄O₆

76. Which kind of isomerism is exhibited by Octahedral [Co(NH₃)₅Br]Cl?

- 1) Ionization isomerism
- 2) Geometrical isomerism
- 3) Optical isomerism
- 4) All of them



- 77. Which of the following is the correct order of increasing field strength of ligands to form coordination compounds?
 - 1) $CO < CN^{-} < en < NH_3 < edta^{-4}$
 - 2) $S^{-2} < Cl^{-} < SCN^{-} < Br^{-} < I^{-}$
 - 3) $NCS^- < H_2O < C_2O_4^{-2} < OH^- < F^-$
 - 4) SCN-<OH-<NCS-<NH₃<CO
- Sol. (4) It is as per the Spectrochemical Series
- 78. The product X and Z in the following reaction are

+
$$CH_3$$
- CH = $CH_2 \xrightarrow{H^+} X$

$$\xrightarrow{O_2/130^{\circ}C} \mathbf{Y} \xrightarrow{H^{+}/H_2O} \mathbf{+Z}$$

- 1) Isopropyl benzene and acetone
- 2) Cumene peroxide and acetone
- 3) Isopropyl benzene and isopropyl alcohol
- 4) Phenol and acetone

79. Aldol condensation reaction is given by

- 2) HCHO
- 3) CH₃-CH₂-CH₂-CHO
- 4) (CH₂)₂C-CHO

80. The end product 'C' in the following sequence of chemical reaction is

$$CH_{_3}COOH \xrightarrow{\quad Ca(OH)_2 \quad} A \xrightarrow{\quad Distillation \quad} B \xrightarrow{\quad NH_2OH \quad} C$$

- 1) Acetaldehyde Oxime
- 2) Formaldehyde Oxime
- 3) Methyl nitrate
- 4) Acetone Oxime

$$CH_3 = NOH$$

$$CH_3 = C$$



- 81. Reaction by which, Benzaldehyde cannot be prepared:
 - +Zn/Hg and conc. HCl
 - + CrO_2Cl_2 in CS_2 followed by $\text{H}_3\text{O}^\oplus$
 - 1 + $\mathrm{H_{2}}$ in presence of Pd+BaSO $_{4}$
 - $\| + CO + HC \|$ in presence of anhydrous AlC $\|$

Sol. (1)

82. What is the major product of the following reaction?

$$\begin{array}{c} \operatorname{CH_3} \\ \operatorname{CH_3} - \operatorname{CH_2} - \operatorname{N-CH_2} - \operatorname{CH_2} - \operatorname{CH_3} & \xrightarrow{\Delta} \\ \operatorname{CH_3} \\ \operatorname{CH_3} \end{array}$$

- 1) $CH_2 = CH_2$
- 2) CH₂=CH-CH₃
- 3) >CH₂
- 4) CH,-CH=CH-CH,

- 83. At high concentration of soap in water, soap behaves as
 - 1) Molecular colloid
 - 2) Associated colloid
 - 3) Macro molecular colloid
 - 4) Lyophilic colloid

Sol. (2)

- 84. Cellulose is a polymer of
 - 1) β-glucose
- 2) α-glucose
- 3) Fructose
- 4) β-galactose

Sol. (1)

- 85. Dettol is an example of
 - 1) Antiseptic
- 2) Antimalarial
- 3) Antibiotic
- 4) Antifertility drug
- Sol. (1) (It is a fact)

Section 'B'

- 86. The de-Broglie's wavelength of electron present in first Bohr orbit of 'H' atom is
 - 1) $\frac{0.529}{2\pi}$ Å
- 2) $2\pi \times 0.529 \text{ Å}$
- 3) 0.529 Å
- 4) 4 × 0.529 Å
- Sol. (2) $\gamma = \frac{2\pi\gamma}{n^2}$
- 87. Heat of combustion of C₂H₄ is -337 K.Cal. If 5.6 lit O, is used at STP in the combustion. Heat liberated is K Cal
 - 1) 28.08
- 2) 14.04
- 3) 42.06
- 4) 56.16
- Sol. (1) $C_1H_4 + 30_2 \longrightarrow 2CO_2 + 2H_0$

88. For the reactions:

$$2NO + O_2 \rightleftharpoons 2NO_2$$
; K_1

$$4NO + 2Cl_2 \rightleftharpoons 4NOCl_2 K_2$$


$$2NO_2 + Cl_2 \rightleftharpoons 2NOCl + O_2$$
; K_3


Where K_1 , K_2 , K_3 are equilibrium constants then K_2^2 equal to:


- 1) $\sqrt{K_2/K_1}$ 2) $\sqrt{K_1K_2}$
- 3) $\sqrt{K_2} / K_1$ 4) $\frac{1}{K_1 K_2}$
- Sol. (3) i) 2NO+02= 2NO2
 - ii) 4NO+2012 = 4NOCI
 - iii) 2NO2+ 01 = 2NOCI+02
 - egniii = egnii egni




89. Which one of the following correctly represents the variation of electronegativity (EN) with atomic number (Z) of group 13 elements?










90. Most stable carbocation in the following

1)
$$(C_6H_5)_2 \overset{+}{C} H$$

2)
$$(CH_3)_3C^+$$



Sol. (4)

It is more stable du to having dancing resonance

91. In which of the following molecules, π -electron density in ring is minimum?

Sol. (4)

92. Equal moles of benzene and toluene are mixed. The vapour pressure of benzene and toluene in pure state are 700 and 600 mm Hg respectively. The mole fraction of benzene in vapour state is

RCC** RCC**

RCC** RCC**

Sol. (4)
$$y_A = \frac{P_A}{P_T} = \frac{P_A^o \times A}{P_A^o \times A + P_B^o \times B}$$
$$\times A = X_B = \frac{1}{2}$$
$$P_A^o = 700 \text{ mm}$$
$$P_B^o = 600 \text{ mm}$$

93. The half life period of a first order chemical reaction is 6.93 minutes. The time required for the completion of 99% of the chemical reaction will be (log 2 = 0.301)

2) 23.03 min

4) 460.6 min

Sol. (3)
$$K = \frac{2.303}{t} \log \frac{A_0}{A_t}$$

$$\frac{0.693}{t_{1/2}} = \frac{2.303}{t} \log \frac{A_0}{A_t}$$

$$\frac{0.693}{6.93} = \frac{2.303}{t} \log \frac{100}{100-99}$$

$$t = 46.06 \text{ min}$$

94. Which of the following electrolysis will have maximum coagulation value for AgI/Ag* sol?

- 1) Na₂S
- 2) Na₃PO₄
- 3) Na₂SO₄
- 4) NaCl

Sol. (4) Coagulation powers

is minimum for Nacl

so coagulation value
is maximum for Nacl.



95. Match the following columns

Column I

Column II

(Metal ions)

(Magnetic moment BM)

1.
$$\sqrt{35}$$

2.
$$\sqrt{30}$$

3.
$$\sqrt{24}$$

4.
$$\sqrt{15}$$

5.
$$\sqrt{8}$$

$$\mathcal{L} = \sqrt{n(n+2)}$$


 $C_{8}^{3+} \Rightarrow n=3$, $F_{8}^{2+} \Rightarrow n=4$
 $N_{8}^{2+} \Rightarrow n=2$, $M_{8}^{2+} \Rightarrow n=5$

96. The spin only magnetic moment of [MnBr₄]²⁻ is 5.9 B.M. The geometry of complex ion is

- 1) Tetrahedral
- 2) Square planar
- 3) Trigonal bipyramidal
- 4) Octahedral

Sol. (1)
$$Mn^{2+} = 3d5$$

 Sp^3 hybn.



$$\frac{C_2H_5O^-/C_2H_5OH}{\text{Mechanism=y}} \xrightarrow{\textbf{B}} \text{Maior}$$

Choose correct option for mechanism x and y, product A and B.


1)
$$x - S_N 2$$
; $y - S_N 2$; $A - \bigcup_{i=1}^{OH} B - \bigcup_{i=1}^{OCH_3} B$


2)
$$x - S_N 1$$
; $y - S_N 2$; A-OH; B -OCH;


3)
$$x - S_N 1$$
; $y - E_2$; $A - \bigcirc OH$; $B - \bigcirc OH$

4)
$$x - S_N 1$$
; $y - E_1$; $A - \bigcirc OH$; $B - \bigcirc OH$

98. If the following is D-glucose, what will be L-Glucose?







Sol. (1)

99. The major product 'Y' in the following reaction

is
$$\stackrel{\text{Cl}}{\longrightarrow} \stackrel{\text{EtONa}}{\longrightarrow} X \stackrel{\text{HBr}}{\longrightarrow} Y$$



100. The product (major) C is

+
$$CO_2 \xrightarrow{\frac{125^{\circ}}{5atm}}$$
 B(major)

 H^+/H_2O and AC_2O



Section 'C' : Botany

Section-A	$\stackrel{\widetilde{\mathbf{P}}}{=} 107.$ plant body is more differentiated the
101. Out of the more than 105 elements discovered	that of algae[11th NCERT Pg. No.43: Summary]
so far less than are essential and	1) Bryophytes 2) Protozoans
beneficial for normal plant growth and	3) Euglenoids 4) Thallophyte
development. [NCERT 11th, summary page No.205]	* Ans. (1)
1) 17 2) 9	2 108. The sporophytes bear to produ
3) 8 4) 21	
Ans. (4)	spores [11th NCERT Pg. No.43: Summary] 1) Sporophyll 2) Microsporophyll
102. Ammonia produced during N ₂ fixation is	
incorporated into _A_ as the _B_ group.	*3) Macrosporophyll 4) Sporangium Ans. (4)
[NCERT 11th, summary page No.205]	† 109 is a lateral outgrowth of stem develope
1) A - amino acid, B - Imino group	at the node [11th NCERT Pg. No 8
2) A - amino acid, B - amino group	Summary]
3) A - amino acid, B - amide group	1) Leaf, Exogenously 2) Bud, Endogenously
4) A -ureids, B - amide group	3) Leaf, Endogenously 4) Flower, Internode
Ans. (2)	
103. Various ions and water from soil can be	Ans. (1)
transported up to a small height in stem by	110 form the basis of classification
[NCERT 11th, summary page No.192] 1) Osmosis 2) Cyclosis	and identification of flowering plants. [11th NCERT Pg. No.82: Summary]
3) Root pressure 4) Diffusion	
Ans. (3)	1) Morphological Characters 2) Anatomical Characters
104. Gametes are formed in haploid organism	
through [NCERT 11th, summary page No.16]	* 3) Floral Characters 4) Vegetative Characters
1) Meiosis 2) Mitosis	1) 1080111111111111111111111111111111111
3) More than 1 correct 4) Syngamy	້ໍບູ້ <mark>Ans. (3)</mark>
Ans. (2)	111. Curly top virus spreads through a plant v
105. Biological classification of plants and animals	the food conducting or phloem tissue.
was first proposed by on the basis	[11th NCERT Pg. No.64: Unit Introduction]
of simple	2) D. J. Ivonwosky
[11th NCERT Pg. No.27: Summary]	2) Maltan Cretton (1) D. Maltanharani
1) Linn., Anatomical Characters	$\stackrel{*}{\overset{*}{\overset{*}{\overset{*}{\overset{*}{\overset{*}{\overset{*}{\overset{*}{$
2) Aristotle, Morphological Character	
3) Linn., Morphological Characters	112. How many of the following are the ma functions of the tissues [11th NCERT Pg. No.
4) Aristotle, colour of blood	
Ans. (2)	* Summary] A) Assimilation of food
106 are cosmopolitan in distribution	
and these organisms show the most extensive	B) Food storage C) transportation of water minerals ar
[11th NCERT Pg. No.27: Summary]	
1) Fungi, Habitat diversity	photosynthates D) Mechanical support
2) Euglenoids, Metabolic diversity	
3) Bacteria, Metabolic diversity	2) O-1 1 4) All 4
4) Protozoans, Habitat diversity	3) Only 1 4) All 4 Ans. (4)
Ans. (3)	∑ Ans. (4)



113. Fats and oils are		RCC**				
1) Glycerol	2) Glycerides	RCC**				
3) Polysaccharides						
Ans. (2)	i) City to interest	RCC				
114. chemosynthetic pathy	way occurs in the	RCC** RCC**				
[11th NCERT Pg. No.224: S						
1) Grana	2) Stroma	RCC**				
3) Lamellae	4) Matrix	RCC**				
Ans. (2)	·					
115. In the light reaction the	e light energy is absorbed	RCC**				
	ent in the antenna, and	RCC**				
-	orophyll a molecule called ² g. No.224: Summary]					
1) Reaction Center		RCC**				
3) LHC	4) Quiescent center					
Ans. (1)	-, 2	RCC** RCC**				
116. The fate of the pyr	uvate depends on the	CC*				
availability of	_					
[11th NCERT Pg. No.237: S	Summary]	RC				
1) Oxygen, organism		RCC** RCC**				
2) Carbon dioxide, oxy	2) Carbon dioxide, oxygen					
3) Nitrogen, Organism	3) Nitrogen, Organism					
4) Type of fermentatio	n, organism	RCC**				
Ans. (1)						
117. Cells of the sporogen	ous tissue lying in the	RCC				
[12th NCERT Pg. No.39-40:	: Summary]	*.				
1) Center of the micro	sporangium	* RC				
2) Periphery of the bil	obed anther	*CC*				
3) Abaxial surface of o	vules	I **:				
4) Adaxial surface of a	nther	* RC				
Ans. (1)		CC*				
118. The central tissues of in to archaesporial ce 40: Summary]	differentiated ells [12th NCERT Pg. No.39-	RCC** RCC** RCC** RCC** RCC** RCC** RCC**				
1) Sporogenous cells	2) Nucellus	C**				
3) Ovules	4) Middle layer	** R(
Ans. (2)	, J -	RCC				

119. Pollen – pistil interaction involves

[12th NCERT Pg. No.39-40: Summary]

- 1) All events from landing of pollen grain on stigma to entry of pollen grain in to embryo sac
- 2) All events from Dispersion through pollen sac of pollen grain to stigma to entry of pollen grain in to embryo sac
- 3) All events from landing of pollen grain on stigma to entry of pollen tube in to embryo sac
- 4) All events from landing of pollen grain on stigma to double fertilization and triple fusion

Ans. (3)

in _____ character that was not expressed in ____ condition may be expressed again when it becomes homozygous. [12th NCERT Pg.No.92-93: Summary]

- 1) Recessive, Heterozygous
- 2) Dominant, Heterozygous
- 3) Heterozygous, Recessive
- 4) Homozygous, heterozygous

Ans. (1)

121. Select the not incorrect statement

[12th NCERT Pg. No.92-93: Summary]

- 1) Not all characters show true dominance because Some characters show incomplete, and some show co-dominance
- 2) All characters show true dominance because incomplete dominance and codominance does not retain the parental combination in further generation
- 3) Some characters incomplete dominance but codominance and parental type is incomplete for the recessive traits
- 4) If codominance is the main criteria, then the parental combination is as same as incomplete dominance.

Ans. (1)

PCB TEST: 1

(21



122. Identify the not-incorrect

[12th NCERT Pg. No.124: Summary]

- 1) DNA and RNA both function as genetic material, but DNA being chemically and structurally more stable is a better genetic material.
- 2) DNA and RNA both function as genetic material, but RNA being chemically and structurally more stable is a better genetic material.
- 3) Only RNA function as genetic material, because DNA being chemically and structurally more unstable is a better genetic material.
- 4) DNA and RNA both function as genetic material, but DNA being chemically and structurally more unstable is a better genetic material.

Ans. (1)

- 123. The DNA replicates semiconservatively, the process is guided by [12th NCERT Pg. No.124: Summary]
 - 1) Antiparallel polynucleotide chain
 - 2) complementary H-bonding
 - 3) Stacking base pair
 - 4) Phosphodiester bond formation

Ans. (2)

- 124. _____ is/are energetically very expensive processes. [12thNCERT Pg. No.124: Summary]
 - 1) Transcription, Translation
 - 2) Translation, Splicing
 - 3) Replication, Translation
 - 4) Splicing, Transcription

Ans. (1)

- 125. The triple crown of biology received by Ernst Mayr includes: [NCERT 11th page 2, Introduction]
 - 1) Crafoord Prize
 - 2) Balan Prize
 - 3) International prize of biotogy
 - 4) All of these

Ans. (4)

- 126. _____ contribute to elongation growth of plant axes. [NCERT 11th page 253, Summary]
 - 1) Root and shoot apical meristems
 - 2) Intercalary meristems
 - 3) Cambium
 - 4) More than one correct

Ans. (4)

- 127. Certain plants need to be exposed to low temperature so as to hasten flowering later in life. This treatment is known as [NCERT 11th page 253, Summary]
 - 1) Plasticity
- 2) Vernalization
- 3) Photoperiodism
- 4) None of these

Ans. (2

- 128. Which of the following is related with Ramdeo Misra
 - 1) Father of Indian Ecology
 - 2) Formulated 1st Post graduate course in ecoloty
 - 3) Due to his efforts, Govt. of India, established the National committee for environment planning and co-ordination [1972], which in later year paved the way for establishment of ministry of environment and forest [1984]
 - 4) All of these

Ans. (4)

RCC**

RCC**

RCC**

RCC**

129. Evolutionary changes through natural selection, takes place at

[NCERT Summary, Page-238, Para-3rd, Line-1,2]

- 1) Organismic level
- 2) Population level
- 3) Community level
- 4) Blome level

Ans. (2)

130. Which of the following statement/s are related with ecosystem and organism respectively

[NCERT Summary, page-256, Line-1,2]

- a) Structural and functional unit of nature
- b) Unit of ecology
- c) Unit of ecological study
- 1) a and b for ecology and c for
- 2) a for ecology and b,c for
- 3) a, b, c for ecosystem only
- 4) a, b, c for organism only

Ans. (1)

RCC**

RCC**



- 131. Rate of assimilation of food energy by consumers is [NCERT Summary, Page-256, Para-2nd, Line-7,8,9]
 - 1) GPP
 - 2) NPP
 - 3) Secondary productivity
 - 4) Respiratory loss

Ans. (3)

- 132. On earth number of inventery [recorded] species and number of estimated species [Which are waiting to be discovered] are
 - 1) 6 million and 1.5 million
 - 2) 1.5 million and 6 million
 - 3) 2 million and 4 million
 - 4) 6 million and 2 million

Ans. (2)

133. Shape of graph for species richness [Species area relation ship]

[NCERT 12th, Summary, Page-268, 2nd Para Last-2nd Lines]

- 1) Rectangular hyperbola
- 2) Linear
- 3) 'J' shaped
- 4) Sigmoid

Ans. (1)

134. Air pollution primarily results from

[VIMP for NEET] [NCERT Summary 12th, Page-285,Line-3,4]

- 1) Deforestation
- 2) Burning of fossil fuel
- 3) Farting at night
- 4) Burning of wood in rural area

Ans. (2)

- 135. The most common source of pollution of water bodies [VIMP for NEET] [NCERT Summary, 12th, Page-285, Line-5,6,7]
 - 1) Domestic sewage
 - 2) Using chemical fertilisers
 - 3) Chemical pesticides
 - 4) Early moring defecation near the bank of river

Ans. (1)

Section-B

- 136. Excess water removed through tips of leaves of plant [NCERT 11th, summary page No.193]
 - 1) Transpiration
- 2) Transduction
- 3) Guttation
- 4) Girdling process

Ans. (3)

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

- 137. The recessive characters are only expressed in _____ conditions and the characters ____ in heterozygous condition [12th NCERT Pg. No.92-93: Summary]
 - 1) Heterozygous, bend
 - 2) Homozygous, never bend
 - 3) Heterozygous, never bend
 - 4) Homozygous, bend

Ans. (2)

- 138. According to Francis Crick Ph.D. study X-ray Diffraction is used [12th NCERT Pg. No.67-68: Unit introduction]
 - 1) Nucleic acids
 - 2) Polypeptides and Proteins
 - 3) Carbohydrates
 - 4) Nucleic acids and Phosphoric acid

Ans. (2)

- 139. A phenomenon called Apomixis is found in [12thNCERT Pg. No.39-40: Summary]
 - 1) A few gymnosperms and particularly Pinus
 - 2) Some angiosperms and particularly citrus
 - 3) a few angiosperms and some Gymnosperm
 - 4) Some angiosperm particularly grasses

Ans. (4)

- 140. Apart from carbohydrates _____ can also be broken down to yield energy. [11thNCERT Pg. No.237-238: Summary]
 - 1) Fats and Proteins
 - 2) Proteins and Glucose
 - 3) Glucose and organic acids
 - 4) Fats and Glucose

Ans. (1)

RCC**



- 141. After absorbing light, electrons are excited and transferred through "A" and "B" and finally to "C" forming "D" [12th NCERT Pg. No.224: Summary]
 - 1) PSI, PSII, NADPH, NAD+
 - 2) PSII, PSI, NAD+, NADPH
 - 3) NAD+, NADPH, PSII, PSI
 - 4) PSII, NAD+, NADPH, PSI

Ans. (2)

- 142. Proteinaceous enzymes exhibit [11th NCERT Pg. No.159-160: Summary]
 - A) Substrate specificity
 - B) Optimum temperature

C) pH

- 1) Only A
- 2) Only B
- 3) Only C
- 4) Only 3

Ans. (1)

- 143. Classification of the vascular bundles is depending on [11thNCERT Pg. No.98: Summary]
 - 1) Presence of Cambium
 - 2) Location of Xylem and Phloem
 - 3) Formation in life span
 - 4) more correct option

Ans. (4)

- 144. How many of the following are criteria help to differentiate the stems from roots [11th NCERT Pg. No.82: Summary]
 - A) Presence of nodes and internodes
 - B) Multicellular hair
 - C) Positively phototropic nature
 - 1) Only 1
- 2) Only 2
- 3) Only 3
- 4) Only A

Ans. (3)

- 145. Algae are classified into three classes on the basis of [11th NCERT Pg. No.43: Summary]
 - A) Type of pigment
 - B) The type of stored food
 - C) The structure of flagella
 - D) The Storage of food
 - 1) Only 1
- 2) Only 2
- 3) Only 3
- 4) Only A

Ans. (2)

- 146. In ecological study, all of the following levels of bioligical organisation are concerned except [NCERT Summary, Page 238, Line-3,4]
 - 1) Ecosystem
- 2) Organism
- 3) Population
- 4) Community

Ans. (1)

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

147. Different organisms are placed in a food chain (ecosystem) on the basis of

[NCERT Summary, Page-256, Line-7,8]

- 1) Source of nutrition
- 2) Type of nutrition
- 3) Amount of food consumed
- 4) Biomass produced

Ans. (1)

148.It is belived that, community with ___ diversity tend to be ____ variable, ___ productive and ____ resistant to biological invasions

[NCERT summary 12th, Page-268, Para-3rd, Line-1,2,3]

- 1) Less, less, more, more
- 2) High, high, less, more
- 3) High, less, less, more
- 4) High, less, more, more

Ans. (4)

149. Soil Pollution, primarily results from

[NCERT Summary, 12th, Page-285, 1st Para, Last 2 lines]

- 1) Agriculture chemicals
- 2) Pesticides
- 3) Leachates from solid waste deposited over pesticides
- 4) All of these

Ans. (4)

- 150. Mendel's law of independent assortment does not hold true for the genes that were located on the [12th NCERT Pg. No.92-93: Summary]
 - 1) Same chromosomes
 - 2) Different chromosomes
 - 3) Homologous chromosomes
 - 4) Non-homologous chromosomes

Ans. (1)

RCC**



Section 'D': Zoology

RCC**

\mathbf{c}			_
50	ct1	on	_ ^
мч	CLL	UII	-/-

- 151. Generation of adenosine triphosphate in mitochondria is [NCERT 11th, summary page No.140]
 - 1) Photo phosphorylation
 - 2) Oxidative phosphorylation
 - 3) Substrate phosphorylation
 - 4) Pseudo phosphorylation

Ans. (2)

- 152. Single membrane structure containing enzyme for digestion of all types of macromolecules [NCERT 12th, summary page No.140]
 - 1) Ribosome
- 2) Deroxisome
- 3) Lysosome
- 4) Centrosome

Ans. (3)

- 153. The process continues throughout life cycle [NCERT 11th, summary page No.170]
 - 1) zygote formation
- 2) sygamy
- 3) More than 1 correct 4) Cell division

Ans. (4)

- 154. The period of cytoplasmic growth [NCERT 11th, summary page No.171]
 - 1) G₁

3) G₂

4) G₀

Ans. (3)

- 155. The parasitic forms show distinct __(i)__ and __(ii)__. Aschelminthes are __(iii)__ include __(iv)__ as well as __(v)__ roondworms [NCERT 11th page 61, Summary]
 - 1) Suckers, hooks, pseudocoelomates, parasitic, non-parasitic
 - 2) Suckers, hooks, coelomates, parasitic, nonparasitic
 - 3) Parapodia, pseudopodia, coelomates, parasitic, non-parasitic
 - 4) More than one correct

Ans. (1)

- 156. Fishes, amphibians, reptiles are _ [NCERT 11th page 61, Summary]
 - 1) Poikilotherms
- 2) Cold-blooded
- 3) Warm blooded
- 4) More than 1 correct

Ans. (4)

- 157. Somatic hybridiation is a process done [NCERT 12th page 178, Summary]
 - 1) Invivo
- 2) Invitro
- 3) Naturally
- 4) Invitro and artifically

Ans. (4)

- 158. The most accepted definition of biotechnology was given by [NCERT 12th page 253, Introduction]
 - 1) WHO
- 2) ICAR
- 3) EFB
- 4) UNESCO

Ans. (3)

- 159. Large scale production involves use of _____ [NCERT 11th page 253, Summary]
 - 1) Fementors
- 2) Bioreactors
- 3) Petri plate culture 4) More than 1 correct

Ans. (4)

- 160. The process to purify the protein or organic compound is called [NCERT 11th page 253, Introduction
 - 1) Upstream process
 - 2) Downstream process
 - 3) Marketing strategy
 - 4) Recombinant DNA technology

Ans. (2)

- 161. The major components of our food are ____ [NCERT-257]
 - 1) Carbohydrate, proteins & Vitamins
 - 2) Carbohydrate, fat & proteins
 - 3) Proteins, vitamins & fat
 - 4) Carbohydrate, proteins, fat, vitamins & minerals

Ans. (2)

- 162. Salivary amylase that digests the starch and converts it into____? [NCERT-266]
 - 1) Glucose
- 2) Maltose
- 3) Sucrose
- 4) all

- 163. Oxygen is utilised by the organisms to ___? [NCERT-268]
 - 1) Directly breakdown simple molecules
 - 2) Indirectly break down simple molecules
 - 3) Directly break down of complex molecules
 - 4) Indirectly breakdown of complex molecules
- Ans. (2)

164. A	sets the p	ace of the activities of the	*
		called pacemaker, here	RCC
	fy the 'A'	[NCERT-288]	*
1) A.V	'. Node	2) SA Node	RC
3) Bur	ndle of His's	4) All	;c**
Ans. (2)			* RC
		ped out by each ventricle cle and it is called the ?	RCC**
		[NCERT-288]	;CC
1) Bea	t volume	2) Cardiac output	**
3) Stro	ke volume	4) More than 1 correct	RCC
Ans. (4)			*
166. Anima	als accumulat	te- [NCERT-290]	RC
1) Am	monia, urea a	nd uric acid	*
2) CO	, water and ic	ons.	* R(
3) Mo	re than one co	orrect	CC*
4) Noi	ne		* "
Ans. (3)			RCC
-	•	aquatic amphibians and	*
aquati	c insects are_	in nature.	RC
		[NCERT-290]	, ,
1) ure		2) Uricotelic	* R(
,	inotelic	4) Ammonotelic	*CC
Ans. (4)			
_		agh which two or more	RCC**
_	s interact and e another.	complemnt the functions [NCERT-315]	
	gration	2) Coordiantion	RC
,	lex action	4) All	, ;
Ans. (2)	ich detion	1) 1 m	* RC
	v important	part of fore braincalled	CC.
		ols the - [NCERT-327]	**
	nperature	2) Eating	RCC
3) Dri	nking	4) All	* * ()
Ans. (4)	Ü		CC** RCC** RCC** RCC** RCC**
170. Father	of 'Green re	volution' in india is -	*. CC

171. Evolutionary biology is the study of -1) History of life forms on earth 2) Fossils 3) Change on earth 4) None Ans. (1) 172. Which of the following provides evidence for evolution -[NCERT-142] 1) Study of comparative anatomy 2) Fossils 3) Comparative biochemistry 4) All Ans. (4) 173. Health is affected by -[NCERT-145] 1) Genetic disorders 2) Infections 3) Life style 4) All Ans. (4) 174. Vector borne disease is -[NCERT-163] 1) Cancer 2) HIV 3) Malaria 4) Cirrhosis Ans. (3) 175. Biology is the youngest of the formalised

disciplines of ____ [NCERT-143]

1) Hypothesis

2) Natural science

3) space science

4) Plaeontology

Ans. (2)

176. In animal body different organ are formed but by using only four types of tissue. It is possible because of [NCERT 11th, Page 100, 2nd Para, Line 2,3,4]

1) Due to modification of cells every time during organ formation

2) Due to arrangement of tissue in a specific proportion and pattern

3) Due to different cell in the same tissue

4) None of these

Ans. (2)

177. How many salivary glands are present in cockroach, near crop

[NCERT 11th, Page 121, Summary 2nd para, line 9,10]

1) One pair

2) Two pair

3) Three pair

4) Three

* 3) T

Ans. (1)

1) M. S. swaminathan

2) James watson

3) Alfonso corti

4) Landsteiner

RCC**

RCC**

[NCERT-144]

178. Alfanso corti was

[NCERT 11th, Page 256, Unit Introduction]

- 1) Italian anatomist
- 2) Americal biologist
- 3) French Naturalist
- 4) Russsian scientist

Ans. (1)

- 179. "Basilar membrane of cochlea contains hair cells, that convert sound vibrations into nerve impulse". This concept was given by
 - 1) Alfanso corti
- 2) Lewis
- 3) Luise pasture
- 4) Rober Brown

Ans. (1)

180. Which of the following statements is/are true

- 1) A female reproductive sytem consists of a pair of ovaries, a pair of oviducts, a uterus, a vagina, external genitalia and a pair of mammary gland
 - [NCERT summary Page-55m, 2nd para line 1,2,3,4]
- 2) Ovarian follicles in different stage of development are embeded in the stroma [NCERT summary, Page-55, 2nd para, line 4,5]
- 3) The process of formation of mature female gamete is oogenesis
- 4) All of these

Ans. (4)

- 181. Mammary glands undergoes differentiation during [NCERT Summary, page 55, Last 3 lines]
 - 1) Pregnancy only
- 2) Puberty
- 3) Menopause
- 4) 1 to 12 years

Ans. (1)

182. Infertility is inability to conseive or produce child, even after _"A"_ years of __B__ sexual cohabitation. A and B respectively are

[NCERT summary, page-65,Last Para,line 1,2]

- 1) One year and unprotected
- 2) Two year and protected
- 3) One year and protected
- 4) Two year and unprotected

Ans. (4)

183. Which of the following country in the world, to initiate various action plans at National level, towards attening a reproductively healthy society

[Most Imp, NCERT Summary, Page-65, Line-2,3,4]

- 1) India
- 2) China
- 3) Russia
- 4) America

Ans. (1)

184. The concept of crop cafeteria, crop scheduling and genetically improving the yield and quality was given by

[NCERT 'Unit Summary' page-144 Para 1 & 2nd]

- 1) Khurana
- 2) Mendel
- 3) Monkambu Sambashivan Swaminathan
- 4) Humbolt

Ans. (3)

RCC**

- 185. The concept of "Lab to land", food security etc. were given by
 - 1) Khurana
 - 2) Mendel
 - 3) Monkambu Sambashivan Swaminathan
 - 4) Humbolt

Ans. (3)

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

RCC**

Section-B

- 186. Select the incorrect statements [NCERT 11th, summary page No.140]
 - 1) The endomembrane system includes ER, golgi complex lysosomes and vacuoles.
 - 2) Centrosome and centriole form the basal body of cilia and flagella
 - 3) Centrioles form spindle apparatus during cell division in all types of cell.
 - 4) Chromoplasts may contain carotene and xanthophyll

- 187. _____ are absent in snakes [NCERT 11th page
 - 61, Summary]
 - 1) Eyes
- 2) Limbs
- 3) parapodia
- 4) water vascolar system

Ans. (2)

- 188. Several new techniques like ____ and ____
 play pivotal role in enhancing food
 production [NCERT 12th page 253, Introduction]
 - 1) Plant tissue culture, animal tissue culture
 - 2) Embryo transfer technology, plant breeding
 - 3) Embryo transfer technology, tissue culture
 - 4) More than one correct

Ans. (3)



189.	Modern bio	technolog	y uses the construction	*
	of	[NCERT 11t	h page 253, Introduction]	RCC
	1) Recombin	nant DNA	2) Chimeric DNA	RCC**
	3) Passenge	r DNA	4) All of these	RC

- 1) Recombinant DNA 2) Chimeric DNA
- 3) Passenger DNA
- 4) All of these

Ans. (4)

190. The undigested food becomes ___ in nature and then enters into the ____? [NCERT-267]

- 1) Solid, rectum
- 2) semisolid, Colon
- 3) Liquify, colon
- 4) Semisolid, Rectum

Ans. (4)

191. Who began his scientific carrer studying the cardiovascular system of reptile and later be turned his attention to the mammalian auditory system? [NCERT-256]

- 1) Alfonso corti
- 2) Ernest mayr
- 4) Eatherine esau
- 4) Melvin colvin

Ans. (1)

192. Creating pressure gradients between the atomsphere and the alveoli with the help of -

[NCERT-276]

- 1) Intercostal muscles and diaphragm
- 2) Diaphraghm and abdominal muscles.
- 3) Intercostal muscles and abdomind muscles
- 4) None

Ans. (1)

193. The pulmonary circulation starts by the blood by the pumping of which is carried to the where it is \mathbf{C} returned to the left atrium. identify the A, B, C, and D. [NCERT-289]

	A	В	С	D
1	Deoxygeneted	Right ventricle	Lungs	Oxygeneted
2	Right ventricle	Lungs	Oxygeneted	Deoxygeneted
3	Lungs	Deoxygeneted	Right ventricle	Oxygeneted
4	Deoxygeneted	Oxygeneted	Lungs	Right ventricle

Ans. (1)

194. Dialysis fluid contain all the constituents as in plasma except___? [NCERT-299]

- 1) Anticoagulant
- 2) Nitrogenous waste.
- 3) More than 1 correct 4) Minerals.

Ans. (2)

195. Point to point rapid coordination among organs provides -[NCERT-331]

- 1) Neural system
- 2) endocrine system
- 3) Integrative system 4) All

Ans. (1)

RCC**

196. Homology is accounted for by the idea of -[NCERT-142]

- 1) Branching descent
- 2) Convergent evolution
- 3) Mutation
- 4) Migration

Ans. (1)

197. How disproved the 'good humor' hypothesis of health. [NCERT-145]

- 1) Discovery of blood circulation
- 2) Demonstration of normal body temperature in presons with black bile
- 3) More than one correct
- 4) Not disproved

Ans. (3)

198. The type of joint which allows "considerable movement" during movement in human body

- is [NCERT 11th, Page-313, Summary, Last two lines]
- 1) Cartilagenous joint 2) Fibrous joint
- 3) Synovial joint
- 4) None of these

Ans. (3)

199. The complex neunendocrine mechanism in parturation involves

- a) Oxytocin
- b) Oestrogen
- c) Cortisol

[NCERT 12th, Page-55, Summary: last 5 lines]

- 1) a, b, c
- 2) a, b
- 3) b, c
- 4) a only

Ans. (1)

RCC**

RCC**

RCC**

200. Antibiotics are used to control all of the following conditions except

[NCERT Summary 12th, Page 188, Last 2 lines]

- 1) Diphtheria
- 2) Whooping cough
- 3) Myocardial infration
- 4) Pneumonia
- Ans. (3)